

Descent of line bundles to GIT quotients of flag varieties by maximal torus

Shrawan Kumar

1 Introduction

Let G be a connected, simply-connected semisimple complex algebraic group with a maximal torus T and let P be a parabolic subgroup containing T . We denote their Lie algebras by the corresponding Gothic characters. The following theorem is one of our main results.

Theorem 3.9 *Let $\mathcal{L}_P(\lambda)$ be a homogeneous ample line bundle on the flag variety $Y = G/P$. Then, the line bundle $\mathcal{L}_P(\lambda)$ descends to a line bundle on the GIT quotient $Y(\lambda)//T$ if and only if for all the semisimple subalgebras $\mathfrak{s}$ of $\mathfrak{g}$ containing $\mathfrak{t}$ (in particular, $\text{rank } \mathfrak{s} = \text{rank } \mathfrak{g}$),*

$$\lambda \in \sum_{\alpha \in \Delta^+(\mathfrak{s})} \mathbb{Z}\alpha,$$

where $\Delta^+(\mathfrak{s})$ is the set of positive roots of $\mathfrak{s}$.

Using the above theorem, we explicitly get exactly which line bundles $\mathcal{L}_P(\lambda)$ descend to $Y(\lambda)//T$. This is our second main result.

In the following Q (resp., Λ) is the root (resp., weight) lattice and we follow the indexing convention as in Bourbaki [B].

Theorem 3.10 *With the notation as in the above theorem, the line bundle $\mathcal{L}_P(\lambda)$ descends to a line bundle on the GIT quotient $Y(\lambda)//T$ if and only if λ is of the following form depending upon the type of G .*

- a) G of type A_ℓ ($\ell \geq 1$) : $\lambda \in Q$.
- b) G of type B_ℓ ($\ell \geq 3$) : $\lambda \in 2Q$.
- c) G of type C_ℓ ($\ell \geq 2$) : $\lambda \in \mathbb{Z}2\alpha_1 + \cdots + \mathbb{Z}2\alpha_{\ell-1} + \mathbb{Z}\alpha_\ell = 2\Lambda$.

- d₁) G of type D_4 : $\lambda \in \{n_1\alpha_1 + 2n_2\alpha_2 + n_3\alpha_3 + n_4\alpha_4 : n_i \in \mathbb{Z} \text{ and } n_1 + n_3 + n_4 \text{ is even}\}$.
- d₂) G of type D_ℓ ($\ell \geq 5$) : $\lambda \in \{2n_1\alpha_1 + 2n_2\alpha_2 + \cdots + 2n_{\ell-2}\alpha_{\ell-2} + n_{\ell-1}\alpha_{\ell-1} + n_\ell\alpha_\ell, n_i \in \mathbb{Z} \text{ and } n_{\ell-1} + n_\ell \text{ is even}\}$.
- e) G of type G_2 : $\lambda \in \mathbb{Z}6\alpha_1 + \mathbb{Z}2\alpha_2$.
- f) G of type F_4 : $\lambda \in \mathbb{Z}6\alpha_1 + \mathbb{Z}6\alpha_2 + \mathbb{Z}12\alpha_3 + \mathbb{Z}12\alpha_4$.
- g) G of type E_6 : $\lambda \in 6\Lambda$.
- h) G of type E_7 : $\lambda \in 12\Lambda$.
- i) G of type E_8 : $\lambda \in 60Q$.

Acknowledgement: We acknowledge partial support from the NSF.

2 Notation

Let G be a connected, simply-connected, semisimple algebraic group over the field $\mathbb{C}$ of complex numbers. We fix a Borel subgroup B of G and a maximal torus T contained in B . Let B^- be the opposite Borel subgroup of G (i.e., the Borel subgroup B^- such that $B^- \cap B = T$). We denote the unipotent radicals of B, B^- by U, U^- respectively. We denote the Lie algebra of any algebraic group by the corresponding Gothic character. In particular, the Lie algebras of G, B, T are denoted by $\mathfrak{g}, \mathfrak{b}, \mathfrak{t}$ respectively. Let $\Delta = \Delta(\mathfrak{g})$ be the set of roots of $\mathfrak{g}$ with respect to $\mathfrak{t}$ and let $\Delta^+ = \Delta^+(\mathfrak{g})$ be the set of positive roots (i.e., the set of roots of $\mathfrak{b}$) with $\Pi = \{\alpha_1, \dots, \alpha_\ell\}$ the set of simple roots. Let W be the Weyl group of G with $\{s_1, \dots, s_\ell\} \subset W$ the set of simple reflections corresponding to the simple roots $\{\alpha_1, \dots, \alpha_\ell\}$ respectively. Let $Q := \bigoplus_{i=1}^\ell \mathbb{Z}\alpha_i \subset \mathfrak{t}^*$ be the root lattice and let

$$\Lambda := \{\lambda \in \mathfrak{t}^* : \lambda(\alpha_i^\vee) \in \mathbb{Z} \text{ for all } i\}$$

be the weight lattice, where $\{\alpha_1^\vee, \dots, \alpha_\ell^\vee\}$ are the simple coroots corresponding to the simple roots $\{\alpha_1, \dots, \alpha_\ell\}$ respectively. Let $\Lambda^+ \subset \Lambda$ be the set of dominant weights, i.e.,

$$\Lambda^+ := \{\lambda \in \mathfrak{t}^* : \lambda(\alpha_i^\vee) \in \mathbb{Z}^+ \text{ for all } i\},$$

where $\mathbb{Z}^+$ is the set of nonnegative integers. For $\lambda \in \Lambda^+$, we denote by $V(\lambda)$ the irreducible representation of $\mathfrak{g}$ with highest weight λ .

Let X be the group of characters of T . Then, X can be identified with Λ by taking the derivative of characters. For $\lambda \in \Lambda$, we denote the corresponding character of T by e^λ . Then, e^λ uniquely extends to a character of B .

We also consider a (standard) parabolic subgroup $P \supset B$ of G . Define

$$\Pi_P = \{\text{simple roots } \alpha_i : -\alpha_i \text{ is a root of } \mathfrak{p}\},$$

$$\Lambda_P = \{\lambda \in \Lambda : \lambda(\alpha_i^\vee) = 0 \text{ for } \alpha_i \in \Pi_P\}, \quad \Lambda_P^+ = \Lambda_P \cap \Lambda^+,$$

and

$$\Lambda_P^o = \{\lambda \in \Lambda_P : \lambda(\alpha_i^\vee) > 0 \text{ for each } \alpha_i \in \Pi \setminus \Pi_P\}.$$

In particular, for $P = B$, $\Lambda_B = \Lambda$ and $\Lambda_B^+ = \Lambda^+$. We recall that e^λ extends to a character of P iff $\lambda \in \Lambda_P$. For $\lambda \in \Lambda_P$, let $\mathcal{L}_P(\lambda) = G \times_P \mathbb{C}_{-\lambda}$ be the homogeneous line bundle on the flag variety G/P associated to the principal P -bundle $G \rightarrow G/P$ via the one dimensional representation $\mathbb{C}_{-\lambda}$ of P given by the character $e^{-\lambda}$. Then, $\mathcal{L}_P(\lambda)$ is an ample line bundle iff $\lambda \in \Lambda_P^o$. For $g \in G$ and $v \in \mathbb{C}_{-\lambda}$, we denote the equivalence class of (g, v) in $G \times_P \mathbb{C}_{-\lambda}$ by $[g, v]$.

3 Proofs

For any set of positive roots $\beta = \{\beta_1, \dots, \beta_n\}$, let $\mathbb{Z}(\beta) := \mathbb{Z}\beta_1 + \dots + \mathbb{Z}\beta_n \subset Q$ and let $\mathfrak{g}(\beta)$ be the semisimple subalgebra of $\mathfrak{g}$ with roots

$$\Delta(\mathfrak{g}(\beta)) := \Delta(\mathfrak{g}) \cap \mathbb{Z}(\beta).$$

We recall the following well-known result.

Lemma 3.1 *For any $\lambda \in \Lambda^+$ and any set of positive roots β such that the index of $\mathbb{Z}(\beta)$ in the root lattice Q is finite, the 0-weight space of the submodule $U(\mathfrak{g}(\beta)) \cdot v_\lambda^+ \subset V(\lambda)$ is nonzero if and only if $\lambda \in \mathbb{Z}(\beta)$, where v_λ^+ is a nonzero highest weight vector of $V(\lambda)$.*

Observe that since $\mathbb{Z}(\beta)$ is of finite index in Q , the full cartan subalgebra $\mathfrak{t} \subset \mathfrak{g}(\beta)$.

Let P be a standard parabolic subgroup. For the action of T on G/P via the left multiplication, the isotropy subgroup $I_{gP} \subset T$ at any $gP \in G/P$ is clearly given by

$$I_{gP} = T \cap gPg^{-1}.$$

Lemma 3.2 *For $gP \in G/P$ and any $\mu \in \Lambda_P$, the isotropy subgroup I_{gP} acts trivially on the fiber $\mathcal{L}_P(\mu)|_{gP}$ iff*

$$e^\mu|_{P \cap g^{-1}Tg} \equiv 1.$$

Proof. For $t \in I_{gP} = T \cap gPg^{-1}$ and any nonzero vector $v_{-\mu} \in \mathbb{C}_{-\mu}$,

$$\begin{aligned} t[g, v_{-\mu}] &= [tg, v_{-\mu}] \\ &= [gg^{-1}tg, v_{-\mu}] \\ &= [g, (g^{-1}tg)v_{-\mu}], \text{ since } g^{-1}tg \in P \\ &= [g, e^{-\mu}(g^{-1}tg)v_{-\mu}]. \end{aligned}$$

Thus,

$$t[g, v_{-\mu}] = [g, v_{-\mu}] \text{ for } t \in T \cap gPg^{-1} \text{ iff } e^{-\mu}(g^{-1}tg) = 1.$$

This proves the lemma. $\square$

Lemma 3.3 *For any $g = \bar{w}up$, for $\bar{w} \in N(T)$, $u \in U_P^-$ and $p \in P$, we have*

$$(1) \quad P \cap g^{-1}Tg = p^{-1}(T \cap u^{-1}Tu)p,$$

where $N(T)$ is the normalizer of T in G and $U_P^- := U^- \cap P$.

Moreover, in the above if we just assume that $u \in U^-$, then

$$(2) \quad P \cap g^{-1}Tg \supset p^{-1}(T \cap u^{-1}Tu)p.$$

Proof. We first prove (1). We have

$$P \cap g^{-1}Tg = P \cap p^{-1}u^{-1}\bar{w}^{-1}T\bar{w}up,$$

i.e.,

$$(3) \quad P \cap g^{-1}Tg = p^{-1}(P \cap u^{-1}Tu)p.$$

Now, for any $t \in T$, if $u^{-1}tu \in P$, then $u^{-1}tu = t$. To prove this, observe that $u^{-1}tut^{-1} \in P \cap U_P^- = (1)$. Thus, $p^{-1}(P \cap u^{-1}Tu)p \subset p^{-1}Tp$ and hence $P \cap g^{-1}Tg \subset p^{-1}Tp \cap g^{-1}Tg$. The reverse inclusion is, of course, obvious.

The inclusion (2) of course follows from (3). $\square$

For $Y = G/P$ and an ample line bundle $\mathcal{L}_P(\lambda)$ on Y (i.e., $\lambda \in \Lambda_P^\circ$), by $Y^{ss}(\lambda)$, we mean the set of semistable points in Y with respect to the action of T via the left multiplication on Y and T -linearized ample line bundle $\mathcal{L}_P(\lambda)$. Then, as is well known, $Y^{ss}(\lambda) \subset Y$ is a Zariski open subset. Moreover, by Lemma 3.1, for any $n \geq 1$, $V(n\lambda)^T \neq 0$ iff $n\lambda \in Q$. Since Q is of finite index in Λ , we get from the next Lemma 3.4 that

$$Y^{ss}(\lambda) \neq \emptyset.$$

Lemma 3.4 *Let $\lambda \in \Lambda_P^\circ$. Then, $gP \in Y^{ss}(\lambda)$ iff $gv_{n\lambda}^+$ has a nonzero component in the zero weight space for some $n \geq 1$, where $v_{n\lambda}^+$ is a highest weight vector of $V(n\lambda)$.*

Proof. By definition, the point $gP \in Y$ is semistable iff there exists a section $\sigma \in H^0(Y, \mathcal{L}_P(n\lambda))^T$ (for some $n \geq 1$), such that $\sigma(gP) \neq 0$. Consider the isomorphism

$$\chi : V(n\lambda)^* \xrightarrow{\sim} H^0(G/P, \mathcal{L}_P(n\lambda)),$$

where $\chi(f)(gP) = [g, (g^{-1}f)|_{\mathbb{C}v_{n\lambda}^+}]$. Thus,

$$\chi(f)(gP) \neq 0 \Leftrightarrow f(gv_{n\lambda}^+) \neq 0,$$

and hence gP is semistable iff $gv_{n\lambda}^+$ has a nonzero component in the zero weight space. $\square$

Lemma 3.5 *For $gP \in Y^{ss}(\lambda)$,*

$$(1) \quad e^{n\lambda}|_{P \cap g^{-1}Tg} \equiv 1, \text{ for some } n \geq 1.$$

In particular,

$$(2) \quad e^\lambda|_{(P \cap g^{-1}Tg)^o} \equiv 1,$$

where $(P \cap g^{-1}Tg)^o$ is the identity component of $P \cap g^{-1}Tg$.

Proof. By Lemma 3.4, gP is semistable iff for some $n \geq 1$, $[gv_{n\lambda}^+]_0 \neq 0$, where $[gv_{n\lambda}^+]_0$ denotes its component in the zero weight space. For $t \in T$,

$$(3) \quad [tgv_{n\lambda}^+]_0 = [gv_{n\lambda}^+]_0.$$

But, for $t \in T \cap gPg^{-1}$,

$$(4) \quad [tgv_{n\lambda}^+]_0 = [gg^{-1}tgv_{n\lambda}^+]_0 = e^{n\lambda}(g^{-1}tg)[gv_{n\lambda}^+]_0.$$

Combining (3) and (4), we get $e^{n\lambda}(g^{-1}tg) = 1$. This proves (1).

The identity (2) follows immediately from (1) since a connected torus is a divisible group. $\square$

Lemma 3.6 *For any $x \in \mathfrak{u}^-$, let β_x be the subset of Δ^+ such that $x = \sum_{\alpha \in \beta_x} x_{-\alpha}$, where $x_{-\alpha} \in \mathfrak{g}_{-\alpha}$ is nonzero. Then, for $u = \text{Exp } x$,*

$$(1) \quad T \cap u^{-1}Tu = \cap_{\alpha \in \beta_x} \{t \in T : e^\alpha(t) = 1\}.$$

In particular, for a weight $\mu \in \Lambda$, $e^\mu|_{T \cap u^{-1}Tu} \equiv 1$ iff $\mu \in \mathbb{Z}\beta_x$.

Proof. Take $t \in T$ such that $utu^{-1} \in T$. Then,

$$utu^{-1}t^{-1} \in T \cap U^- = \{1\}.$$

Thus, $utu^{-1} = t$, i.e., $tut^{-1} = u$. But,

$$\begin{aligned} tut^{-1} &= \text{Exp}(\text{Ad } t \cdot x) \\ &= \text{Exp}\left(\sum_{\alpha \in \beta_x} e^{-\alpha}(t)x_{-\alpha}\right) \\ &= \text{Exp}(x). \end{aligned}$$

Since $\text{Exp}|_{\mathfrak{u}^-}$ is a bijection,

$$\sum_{\alpha \in \beta_x} e^{-\alpha}(t)x_{-\alpha} = \sum_{\alpha \in \beta_x} x_{-\alpha}.$$

Thus, $e^{-\alpha}(t) = 1$ for any $\alpha \in \beta_x$. This proves the inclusion

$$T \cap u^{-1}Tu \subset \cap_{\alpha \in \beta_x} \{t \in T : e^\alpha(t) = 1\}.$$

The reverse inclusion follows by reversing the above calculation.

To prove the ‘In particular’ statement, consider the isomorphism $\xi : T \simeq \text{Hom}_{\mathbb{Z}}(\Lambda, \mathbb{C}^*)$, $\xi(t)\mu = e^\mu(t)$, for $t \in T$ and $\mu \in \Lambda$. From this identification and (1) it is easy to see that $\xi(T \cap u^{-1}Tu) = \text{Hom}_{\mathbb{Z}}(\frac{\Lambda}{\mathbb{Z}\beta_x}, \mathbb{C}^*)$. Now, if $\mu \in \Lambda \setminus (\mathbb{Z}\beta_x)$, there exists a homomorphism $f : \frac{\Lambda}{\mathbb{Z}\beta_x} \rightarrow \mathbb{C}^*$ Such that $f(\mu) \neq 1$. From this we conclude the ‘In particular’ statement. $\square$

Since $Y^{ss}(\lambda)$ is Zariski open in Y , we can find a semistable point of the form $\text{Exp}(x)P$, where $x \in \mathfrak{u}^-$ and $\beta_x = \Delta^+$.

More generally, take any subset $\beta \subset \Delta^+$ such that $\mathbb{Z}(\beta)$ is of finite index in Q . Then, for some $n \geq 1$, $n\lambda \in \mathbb{Z}(\beta)$ and hence by Lemma 3.1,

$$(U(\mathfrak{g}(\beta)) \cdot v_{n\lambda}^+)^T \neq (0).$$

In particular,

$$(G(\beta)/B(\beta)) \cap Y^{ss}(\lambda) \neq \emptyset,$$

where $G(\beta)$ is the connected (semisimple) subgroup of G with Lie algebra $\mathfrak{g}(\beta)$ and $B(\beta) := B \cap G(\beta)$ is a Borel subgroup of $G(\beta)$. Again, by Zariski density, we can find an element of the form $\text{Exp}(x)P \in Y^{ss}(\lambda)$ such that $x \in \mathfrak{u}^-$ and β_x is the set of all the positive roots of $G(\beta)$ (i.e., all the roots of $B(\beta)$).

Lemma 3.7 *For any subset $S \subset \Delta^+$, the quotient group*

$$T_S/T_S^o \simeq \text{Tor}(\Lambda/\mathbb{Z}S),$$

where $T_S := \cap_{\alpha \in S} \{t \in T : e^\alpha(t) = 1\}$, T_S^o denotes its identity component and Tor denotes the torsion subgroup.

Proof. Recall the identification $\xi : T \xrightarrow{\sim} \text{Hom}_{\mathbb{Z}}(\Lambda, \mathbb{C}^*)$ from the proof of Lemma 3.6. Under this identification

$$\xi(T_S) = \text{Hom}_{\mathbb{Z}}\left(\frac{\Lambda}{\mathbb{Z}S}, \mathbb{C}^*\right).$$

Decompose

$$\frac{\Lambda}{\mathbb{Z}S} \simeq \text{Tor}\left(\frac{\Lambda}{\mathbb{Z}S}\right) \oplus F,$$

where F is a free $\mathbb{Z}$ -module. Thus,

$$\text{Hom}_{\mathbb{Z}}\left(\frac{\Lambda}{\mathbb{Z}S}, \mathbb{C}^*\right) \simeq \text{Hom}_{\mathbb{Z}}\left(\text{Tor}\left(\frac{\Lambda}{\mathbb{Z}S}\right), \mathbb{C}^*\right) \times \text{Hom}_{\mathbb{Z}}(F, \mathbb{C}^*).$$

But, $\mathrm{Hom}_{\mathbb{Z}}(F, \mathbb{C}^*) \simeq (\mathbb{C}^*)^m$, where $m := \mathrm{rk} F$ and, of course,

$$\mathrm{Hom}_{\mathbb{Z}}(\mathrm{Tor}(\frac{\Lambda}{\mathbb{Z}S}, \mathbb{C}^*) \simeq \mathrm{Tor}(\frac{\Lambda}{\mathbb{Z}S}).$$

Thus, $T_S^o \simeq \mathrm{Hom}_{\mathbb{Z}}(F, \mathbb{C}^*)$ and hence

$$T_S/T_S^o \simeq \mathrm{Tor}(\Lambda/\mathbb{Z}S). \quad \square$$

Let H be a reductive group, X a projective variety and $\mathcal{L}$ an ample H -equivariant line bundle on X . Then, recall that the *GIT quotient* $X//H$ is by definition the uniform categorical quotient of the (open) set of semistable points X^{ss} by H (cf. [MFK, Theorem 1.10]).

We recall the following ‘descent’ lemma of Kempf (cf. [DN, Theorem 2.3]).

Lemma 3.8 *Let X, H and $\mathcal{L}$ be as above. Then, an H -equivariant vector bundle $\mathcal{S}$ on X descends to a vector bundle on $X//H$ (i.e., there exists a vector bundle $\mathcal{S}'$ on $X//H$ such that its pull-back to X^{ss} under the canonical H -equivariant structure is H -equivariantly isomorphic to the restriction of $\mathcal{S}$ to X^{ss}) iff for any $x \in X^{ss}$, the isotropy subgroup I_x acts trivially on the fiber $\mathcal{S}_x$.*

In fact (though we do not need), for the ‘if’ part, it suffices to assume that I_x acts trivially for only those $x \in X^{ss}$ such that the orbit $H \cdot x$ is closed in X^{ss} .

Let P be a standard parabolic subgroup and let $\mathcal{L}_P(\lambda)$ be a homogeneous ample line bundle (i.e., $\lambda \in \Lambda_P^o$) on $Y = G/P$. Denote the GIT quotient of Y by T with respect to $\mathcal{L}_P(\lambda)$ by $Y(\lambda)//T$. The following is one of our main results.

Theorem 3.9 *With the notation as above, the line bundle $\mathcal{L}_P(\lambda)$ descends to a line bundle on $Y(\lambda)//T$ iff for all the semisimple subalgebras $\mathfrak{s}$ of $\mathfrak{g}$ containing $\mathfrak{t}$ (in particular, $\mathrm{rank} \mathfrak{s} = \mathrm{rank} \mathfrak{g}$),*

$$\lambda \in \mathbb{Z}\Delta^+(\mathfrak{s}),$$

where $\Delta^+(\mathfrak{s}) := \Delta^+ \cap \Delta(\mathfrak{s})$ is the set of positive roots of $\mathfrak{s}$.

Proof. By Lemma 3.8, $\mathcal{L}_P(\lambda)$ descends to $Y(\lambda)//T$ iff for all the semistable points $gP = \bar{w}uP \in G/P$ (for $\bar{w} \in N(T)$ and $u \in U_P^-$), the isotropy subgroup I_{gP} acts trivially on the fiber $\mathcal{L}_P(\lambda)|_{gP}$. By Lemmas 3.2 and 3.3, this is equivalent to the requirement that $e^\lambda|_{T \cap u^{-1}Tu} \equiv 1$. Further, by Lemma 3.6, this is equivalent to the requirement that $\lambda \in \mathbb{Z}\beta_x$, where $x \in \mathfrak{u}^-$ is the element with $\text{Exp } x = u$.

By the discussion above Lemma 3.7, for any semisimple subalgebra $\mathfrak{s}$ of $\mathfrak{g}$ containing $\mathfrak{t}$, there exists an element $x \in \mathfrak{u}^-$ such that $\text{Exp}(x)P \in G/P$ is semistable and, moreover, $\beta_x = \Delta^+(\mathfrak{s})$. Thus, we get by Lemmas 3.2, 3.3 and 3.6 that $\lambda \in \mathbb{Z}\Delta^+(\mathfrak{s})$ for any such $\mathfrak{s}$ if the line bundle $\mathcal{L}_P(\lambda)$ descends to $Y(\lambda)//T$.

Conversely, take any semistable point $\bar{w}\text{Exp}(x)P \in G/P$ for $\bar{w} \in N(T)$ and $x \in \mathfrak{u}_P^-$. If $\mathbb{Z}\beta_x \subset Q$ is not of finite index in Q , choose simple roots $\alpha_x = \{\alpha_{i_1}, \dots, \alpha_{i_j}\}$ such that $\mathbb{Q}\beta_x \cap \mathbb{Q}\alpha_x = (0)$ and $\mathbb{Q}\alpha_x + \mathbb{Q}\beta_x = \bigoplus_{i=1}^\ell \mathbb{Q}\alpha_i$, where $\mathbb{Q}\beta_x := \sum_{\beta \in \beta_x} \mathbb{Q}\beta \subset \mathfrak{t}^*$ and $\mathbb{Q}\alpha_x := \bigoplus_{n=1}^j \mathbb{Q}\alpha_{i_n} \subset \mathfrak{t}^*$. With this choice of α_x , the torsion submodule

$$(1) \quad \text{Tor}(\Lambda/\mathbb{Z}\beta_x) \hookrightarrow \text{Tor}(\Lambda/(\mathbb{Z}\beta_x + \mathbb{Z}\alpha_x)).$$

To see this observe that we have the following short exact sequence:

$$0 \rightarrow \mathbb{Z}\alpha_x \rightarrow \Lambda/\mathbb{Z}\beta_x \rightarrow \Lambda/(\mathbb{Z}\alpha_x + \mathbb{Z}\beta_x) \rightarrow 0.$$

From this the assertion (1) follows easily.

Let $\mathfrak{s}$ be the semisimple subalgebra of $\mathfrak{g}$ containing $\mathfrak{t}$ such that

$$\mathbb{Z}\Delta^+(\mathfrak{s}) = \mathbb{Z}\beta_x + \mathbb{Z}\alpha_x.$$

Choose an element $y = y_{\mathfrak{s}} \in \mathfrak{u}^-$ such that $\beta_y = \Delta^+(\mathfrak{s})$ and $(\text{Exp } y)P \in Y^{ss}(\lambda)$. With this choice of y , by Lemma 3.6, we see that $T \cap v^{-1}Tv \subset T \cap u^{-1}Tu$, where $u := \text{Exp } x$ and $v := \text{Exp } y$. We further show that the inclusion $T \cap v^{-1}Tv \subset T \cap u^{-1}Tu$ induces a splitting:

$$(2) \quad (T \cap u^{-1}Tu)/(T \cap u^{-1}Tu)^o \hookrightarrow (T \cap v^{-1}Tv)/(T \cap v^{-1}Tv)^o.$$

By Lemma 3.7,

$$(T \cap v^{-1}Tv)/(T \cap v^{-1}Tv)^o \simeq \text{Tor}\left(\frac{\Lambda}{\mathbb{Z}\beta_y}\right)$$

and

$$(T \cap u^{-1}Tu)/(T \cap u^{-1}Tu)^o \cong \text{Tor}\left(\frac{\Lambda}{\mathbb{Z}\beta_x}\right).$$

Combining the above identifications with (1), we get (2).

Since $\bar{w}uP$ is a semistable point, by Lemmas 3.3 and 3.5, $e^\lambda|_{(T \cap u^{-1}Tu)^o} = 1$. Moreover, by the assumption and Lemma 3.6, $e^\lambda|_{T \cap v^{-1}Tv} \equiv 1$. Thus, $e^\lambda|_{T \cap u^{-1}Tu} \equiv 1$ by (2). This proves the theorem. $\square$

Using the above theorem, we explicitly get exactly which $\mathcal{L}_P(\lambda)$ descends to the GIT quotient $Y(\lambda)//T$.

In the following, we follow the indexing convention as in Bourbaki [B, Planche I - IX].

Theorem 3.10 *Let G be a connected, simply-connected simple algebraic group, $P \subset G$ a standard parabolic subgroup and let $\mathcal{L}_P(\lambda)$ be a homogeneous ample line bundle on the flag variety $Y = G/P$ (i.e., $\lambda \in \Lambda_P^o$). Then, the line bundle $\mathcal{L}_P(\lambda)$ descends to a line bundle on the GIT quotient $Y(\lambda)//T$ if and only if λ is of the following form depending upon the type of G (in addition to $\lambda \in \Lambda_P^o$).*

a) G of type A_ℓ ($\ell \geq 1$) : $\lambda \in Q$.

b) G of type B_ℓ ($\ell \geq 3$) : $\lambda \in 2Q$.

c) G of type C_ℓ ($\ell \geq 2$) : $\lambda \in \mathbb{Z}2\alpha_1 + \cdots + \mathbb{Z}2\alpha_{\ell-1} + \mathbb{Z}\alpha_\ell = 2\Lambda$.

d₁) G of type D_4 : $\lambda \in \{n_1\alpha_1 + 2n_2\alpha_2 + n_3\alpha_3 + n_4\alpha_4 : n_i \in \mathbb{Z} \text{ and } n_1 + n_3 + n_4 \text{ is even}\}$.

d₂) G of type D_ℓ ($\ell \geq 5$) : $\lambda \in \{2n_1\alpha_1 + 2n_2\alpha_2 + \cdots + 2n_{\ell-2}\alpha_{\ell-2} + n_{\ell-1}\alpha_{\ell-1} + n_\ell\alpha_\ell, n_i \in \mathbb{Z} \text{ and } n_{\ell-1} + n_\ell \text{ is even}\}$.

e) G of type G_2 : $\lambda \in \mathbb{Z}6\alpha_1 + \mathbb{Z}2\alpha_2$.

f) G of type F_4 : $\lambda \in \mathbb{Z}6\alpha_1 + \mathbb{Z}6\alpha_2 + \mathbb{Z}12\alpha_3 + \mathbb{Z}12\alpha_4$.

g) G of type E_6 : $\lambda \in 6\Lambda$.

h) G of type E_7 : $\lambda \in 12\Lambda$.

i) G of type E_8 : $\lambda \in 60Q$.

Proof. The theorem follows by using Theorem 3.9 and the classification of the semisimple subalgebras of $\mathfrak{g}$ of maximal rank due to Borel-Siebenthal (cf. [W, Theorem 8.10.8]). In the following chart, we only list all the proper maximal semisimple subalgebras $\mathfrak{s}$ of $\mathfrak{g}$ containing the (fixed) Cartan subalgebra $\mathfrak{t}$ up to a conjugation under the Weyl group. In the following, θ denotes the highest root and $\mathfrak{s}_i$ denotes the semisimple subalgebra of $\mathfrak{g}$ containing $\mathfrak{t}$ with simple roots $\{\alpha_1, \dots, \widehat{\alpha}_i, \dots, \alpha_\ell, -\theta\}$.

- a) A_ℓ ($\ell \geq 1$): none
- b) B_ℓ ($\ell \geq 3$) : $\mathfrak{s}_i, 2 \leq i \leq \ell$
- c) C_ℓ ($\ell \geq 2$) : $\mathfrak{s}_i, 1 \leq i \leq \ell - 1$
- d) D_ℓ ($\ell \geq 4$) : $\mathfrak{s}_i, 2 \leq i \leq \ell - 2$
- e) G_2 : $\mathfrak{s}_i, i=1,2$
- f) F_4 : $\mathfrak{s}_i, i = 1, 2, 4$
- g) E_6 : $\mathfrak{s}_i, i = 2, 3, 4, 5$
- h) E_7 : $\mathfrak{s}_i, i = 1, 2, 3, 5, 6$
- i) E_8 : $\mathfrak{s}_i, i = 1, 2, 5, 7, 8$.

We denote by $L(\mathfrak{g}; \alpha_1, \dots, \alpha_\ell)$ the intersection of $\mathbb{Z}\Delta^+(\mathfrak{s})$ (inside the root lattice Q) as $\mathfrak{s}$ varies over all possible semisimple subalgebras $\mathfrak{s}$ of $\mathfrak{g}$ containing the (fixed) cartan subalgebra $\mathfrak{t}$.

In the following, we determine $L(\mathfrak{g}; \alpha_1, \dots, \alpha_\ell)$ for each simple $\mathfrak{g}$.

- a) A_ℓ ($\ell \geq 1$) : In this case $L(A_\ell; \alpha_1, \dots, \alpha_\ell) = Q$.
- c) C_ℓ ($\ell \geq 2$) : By the above chart,

$$\begin{aligned} L(C_\ell; \alpha_1, \dots, \alpha_\ell) &= [(\mathbb{Z}\theta + L(C_{\ell-1}; \alpha_2, \dots, \alpha_\ell)) \cap \\ &\quad (L(C_2; \alpha_1, -\theta) + L(C_{\ell-2}; \alpha_3, \dots, \alpha_\ell)) \cap \dots \\ &\quad (L(C_{\ell-2}; \alpha_{\ell-3}, \dots, \alpha_1, -\theta) + L(C_2; \alpha_{\ell-1}, \alpha_\ell)) \cap \\ &\quad (L(C_{\ell-1}; \alpha_{\ell-2}, \dots, \alpha_1, -\theta) + \mathbb{Z}\alpha_\ell)]_W, \end{aligned}$$

where $[M]_W$ denotes $\cap_{w \in W} wM$. By induction, for $j < \ell$,

$$L(C_j; \alpha_{\ell-j+1}, \dots, \alpha_\ell) = \mathbb{Z}2\alpha_{\ell-j+1} + \dots + \mathbb{Z}2\alpha_{\ell-1} + \mathbb{Z}\alpha_\ell$$

and

$$\begin{aligned} L(C_j; \alpha_{j-1}, \dots, \alpha_1, -\theta) &= \mathbb{Z}2\alpha_{j-1} + \dots + \mathbb{Z}2\alpha_1 + \mathbb{Z}\theta \\ &= \mathbb{Z}2\alpha_{j-1} + \dots + \mathbb{Z}2\alpha_1 + \mathbb{Z}(2\alpha_j + \dots + 2\alpha_{\ell-1} + \alpha_\ell), \end{aligned}$$

since

$$\theta(C_\ell; \alpha_1, \dots, \alpha_\ell) = 2\alpha_1 + \dots + 2\alpha_{\ell-1} + \alpha_\ell.$$

Thus,

$$L(C_\ell; \alpha_1, \dots, \alpha_\ell) = [\mathbb{Z}2\alpha_1 + \dots + \mathbb{Z}2\alpha_{\ell-1} + \mathbb{Z}\alpha_\ell]_W.$$

But, $\mathbb{Z}2\alpha_1 + \dots + \mathbb{Z}2\alpha_{\ell-1} + \mathbb{Z}\alpha_\ell$ is W -stable and hence

$$L(C_\ell; \alpha_1, \dots, \alpha_\ell) = \mathbb{Z}2\alpha_1 + \dots + \mathbb{Z}2\alpha_{\ell-1} + \mathbb{Z}\alpha_\ell.$$

This proves the (c)-part of the theorem.

d) D_ℓ ($\ell \geq 4$) : We first consider the case of D_4 . In this case

$$\begin{aligned} L(D_4; \alpha_1, \dots, \alpha_4) &= [L(A_1; \alpha_1) + L(A_1; -\theta) + L(A_1; \alpha_3) + L(A_1; \alpha_4)]_W \\ &= [\mathbb{Z}\alpha_1 + \mathbb{Z}\theta + \mathbb{Z}\alpha_3 + \mathbb{Z}\alpha_4]_W \\ &= [\mathbb{Z}\alpha_1 + \mathbb{Z}2\alpha_2 + \mathbb{Z}\alpha_3 + \mathbb{Z}\alpha_4]_W. \end{aligned}$$

Now, the sublattice $L' := \{n_1\alpha_1 + n_2\alpha_2 + n_3\alpha_3 + n_4\alpha_4 : n_i \in \mathbb{Z}, n_2 \text{ is even and } \sum_{i=1}^4 n_i \text{ is even}\} \subset \mathbb{Z}\alpha_1 + \mathbb{Z}2\alpha_2 + \mathbb{Z}\alpha_3 + \mathbb{Z}\alpha_4$ is W -stable (as is easy to see). Moreover, clearly the index of L' in $\mathbb{Z}\alpha_1 + \mathbb{Z}2\alpha_2 + \mathbb{Z}\alpha_3 + \mathbb{Z}\alpha_4$ is 2 and $\mathbb{Z}\alpha_1 + \mathbb{Z}2\alpha_2 + \mathbb{Z}\alpha_3 + \mathbb{Z}\alpha_4$ is not W -stable (as can be seen by applying the second simple reflection s_2). Thus $[\mathbb{Z}\alpha_1 + \mathbb{Z}2\alpha_2 + \mathbb{Z}\alpha_3 + \mathbb{Z}\alpha_4]_W = L'$, proving the theorem in this case.

We now come to the case of general D_ℓ . By the above chart and the A_ℓ case,

$$\begin{aligned} L(D_\ell; \alpha_1, \dots, \alpha_\ell) &= [(L(D_2; \alpha_1, -\theta) + L(D_{\ell-2}; \alpha_3, \dots, \alpha_\ell)) \cap \dots \\ &\quad \cap (L(D_{\ell-2}; \alpha_{\ell-3}, \dots, \alpha_1, -\theta) + L(D_2; \alpha_{\ell-1}, \alpha_\ell))]_W, \end{aligned}$$

where D_k for $k = 2, 3$ is interpreted as A_k . Set

$$L'(\alpha_{\ell-3}, \alpha_{\ell-2}, \alpha_{\ell-1}, \alpha_\ell) = \left\{ \sum_{i=\ell-3}^{\ell} n_i \alpha_i : n_i \in \mathbb{Z}, \sum n_i \text{ is even and } n_{\ell-2} \text{ is even} \right\},$$

and if $\ell - k > 4$, set

$$L'(\alpha_{k+1}, \dots, \alpha_\ell) = \left\{ \sum_{i=k+1}^{\ell} n_i \alpha_i : n_i \in \mathbb{Z}, \sum n_i \text{ is even and } n_{k+1}, \dots, n_{\ell-2} \text{ are even} \right\},$$

and if $\ell - k < 4$, set

$$L'(\alpha_{k+1}, \dots, \alpha_\ell) = \mathbb{Z}\alpha_{k+1} + \dots + \mathbb{Z}\alpha_\ell.$$

By induction and the $A_{\ell-k}$ case, for $2 \leq \ell - k < \ell$,

$$L(D_{\ell-k}; \alpha_{k+1}, \dots, \alpha_\ell) = L'(\alpha_{k+1}, \dots, \alpha_\ell).$$

Thus,

$$\begin{aligned} L(D_\ell; \alpha_1, \dots, \alpha_\ell) &= [(\mathbb{Z}\alpha_1 + \mathbb{Z}\theta + L'(\alpha_3, \dots, \alpha_\ell)) \cap \\ &\quad (\mathbb{Z}\alpha_1 + \mathbb{Z}\alpha_2 + \mathbb{Z}\theta + L'(\alpha_4, \dots, \alpha_\ell)) \cap \\ &\quad (L'(\alpha_3, \alpha_2, \alpha_1, -\theta) + L'(\alpha_5, \dots, \alpha_\ell)) \cap \dots \\ &\quad \cap (L'(\alpha_{\ell-5}, \dots, \alpha_1, -\theta) + L'(\alpha_{\ell-3}, \alpha_{\ell-2}, \alpha_{\ell-1}, \alpha_\ell)) \\ &\quad \cap (L'(\alpha_{\ell-4}, \dots, \alpha_1, -\theta) + \mathbb{Z}\alpha_{\ell-2} + \mathbb{Z}\alpha_{\ell-1} + \mathbb{Z}\alpha_\ell) \\ &\quad \cap (L'(\alpha_{\ell-3}, \dots, \alpha_1, -\theta) + \mathbb{Z}\alpha_{\ell-1} + \mathbb{Z}\alpha_\ell)]_W \\ &= [L'(\alpha_1, \dots, \alpha_\ell)]_W. \end{aligned}$$

It is easy to see that $L'(\alpha_1, \dots, \alpha_\ell)$ is W -invariant and hence

$$[L'(\alpha_1, \dots, \alpha_\ell)]_W = L'(\alpha_1, \dots, \alpha_\ell).$$

This proves the theorem in the case of D_ℓ .

(b) B_ℓ ($\ell \geq 3$): By the above chart, we get that

$$\begin{aligned} L(B_\ell; \alpha_1, \dots, \alpha_\ell) &= [(\mathbb{Z}\alpha_1 + \mathbb{Z}\theta + L(B_{\ell-2}; \alpha_3, \dots, \alpha_\ell)) \cap \\ &\quad (L(A_3; \alpha_1, \alpha_2, -\theta) + L(B_{\ell-3}; \alpha_4, \dots, \alpha_\ell)) \cap \\ &\quad (L(D_4; \alpha_3, \alpha_2, \alpha_1, -\theta) + L(B_{\ell-4}; \alpha_5, \dots, \alpha_\ell)) \cap \dots \\ &\quad (L(D_{\ell-3}; \alpha_{\ell-4}, \dots, \alpha_1, -\theta) + L(B_3; \alpha_{\ell-2}, \alpha_{\ell-1}, \alpha_\ell)) \cap \\ &\quad (L(D_{\ell-2}; \alpha_{\ell-3}, \dots, \alpha_1, -\theta) + L(C_2; \alpha_\ell, \alpha_{\ell-1})) \cap \\ &\quad (L(D_{\ell-1}; \alpha_{\ell-2}, \dots, \alpha_1, -\theta) + \mathbb{Z}\alpha_\ell) \cap \\ &\quad L(D_\ell; \alpha_{\ell-1}, \dots, \alpha_1, -\theta)]_W. \end{aligned}$$

By using the result for D_ℓ and A_3 and also, by induction, for B_j ($j < \ell$), we get that (with L' as in the proof of the D_ℓ case)

$$\begin{aligned} L(B_\ell; \alpha_1, \dots, \alpha_\ell) &= [(\mathbb{Z}\alpha_1 + \mathbb{Z}\theta + \mathbb{Z}2\alpha_3 + \dots + \mathbb{Z}2\alpha_\ell) \cap \\ &\quad (\mathbb{Z}\alpha_1 + \mathbb{Z}\alpha_2 + \mathbb{Z}\theta + \mathbb{Z}2\alpha_4 + \dots + \mathbb{Z}2\alpha_\ell) \cap \\ &\quad (L'(\alpha_3, \alpha_2, \alpha_1, -\theta) + \mathbb{Z}2\alpha_5 + \dots + \mathbb{Z}2\alpha_\ell) \cap \dots \\ &\quad (L'(\alpha_{\ell-4}, \dots, \alpha_1, -\theta) + \mathbb{Z}2\alpha_{\ell-2} + \mathbb{Z}2\alpha_{\ell-1} + \mathbb{Z}2\alpha_\ell) \cap \\ &\quad (L'(\alpha_{\ell-3}, \dots, \alpha_1, -\theta) + \mathbb{Z}2\alpha_\ell + \mathbb{Z}\alpha_{\ell-1}) \cap \\ &\quad (L'(\alpha_{\ell-2}, \dots, \alpha_1, -\theta) + \mathbb{Z}\alpha_\ell) \cap L'(\alpha_{\ell-1}, \dots, \alpha_1, -\theta)]_W \\ &= [\mathbb{Z}2\alpha_1 + \dots + \mathbb{Z}2\alpha_\ell]_W \\ &= 2[Q]_W \\ &= 2Q, \text{ since } Q \text{ is } W\text{-stable.} \end{aligned}$$

This proves the theorem for the case of B_ℓ .

(e) G_2 : By the chart and the theorem for the case of A_2 ,

$$\begin{aligned}
L(G_2; \alpha_1, \alpha_2) &= [(L(A_2; \alpha_2, -\theta)) \cap (\mathbb{Z}\alpha_1 + \mathbb{Z}\theta)]_W \\
&= [(\mathbb{Z}\alpha_2 + \mathbb{Z}\theta) \cap (\mathbb{Z}\alpha_1 + \mathbb{Z}\theta)]_W \\
&= [(\mathbb{Z}\alpha_2 + \mathbb{Z}3\alpha_1) \cap (\mathbb{Z}\alpha_1 + \mathbb{Z}2\alpha_2)]_W \\
&= [\mathbb{Z}3\alpha_1 + \mathbb{Z}2\alpha_2]_W \\
&= \mathbb{Z}6\alpha_1 + \mathbb{Z}2\alpha_2,
\end{aligned}$$

since $\mathbb{Z}6\alpha_1 + \mathbb{Z}2\alpha_2$ is W -stable, whereas $\mathbb{Z}3\alpha_1 + \mathbb{Z}2\alpha_2$ is not W -stable and $\mathbb{Z}6\alpha_1 + \mathbb{Z}2\alpha_2$ is of index 2 inside $\mathbb{Z}3\alpha_1 + \mathbb{Z}2\alpha_2$.

(f) F_4 : By the chart,

$$\begin{aligned}
L(F_4; \alpha_1, \alpha_2, \alpha_3, \alpha_4) &= [(\mathbb{Z}\theta + L(C_3; \alpha_4, \alpha_3, \alpha_2)) \cap \\
&\quad (L(A_2; -\theta, \alpha_1) + L(A_2; \alpha_3, \alpha_4)) \cap \\
&\quad L(B_4; -\theta, \alpha_1, \alpha_2, \alpha_3)]_W
\end{aligned}$$

By the theorem for A_2, C_3 and B_4 , we get

$$\begin{aligned}
L(F_4; \alpha_1, \alpha_2, \alpha_3, \alpha_4) &= [(\mathbb{Z}\theta + \mathbb{Z}2\alpha_4 + \mathbb{Z}2\alpha_3 + \mathbb{Z}\alpha_2) \cap \\
&\quad (\mathbb{Z}\theta + \mathbb{Z}\alpha_1 + \mathbb{Z}\alpha_3 + \mathbb{Z}\alpha_4) \cap \\
&\quad (\mathbb{Z}2\theta + \mathbb{Z}2\alpha_1 + \mathbb{Z}2\alpha_2 + \mathbb{Z}2\alpha_3)]_W \\
&= [(\mathbb{Z}2\alpha_1 + \mathbb{Z}2\alpha_4 + \mathbb{Z}2\alpha_3 + \mathbb{Z}\alpha_2) \cap \\
&\quad (\mathbb{Z}3\alpha_2 + \mathbb{Z}\alpha_1 + \mathbb{Z}\alpha_3 + \mathbb{Z}\alpha_4) \cap \\
&\quad (\mathbb{Z}4\alpha_4 + \mathbb{Z}2\alpha_1 + \mathbb{Z}2\alpha_2 + \mathbb{Z}2\alpha_3)]_W \\
&= [M]_W, \text{ where } M := \mathbb{Z}2\alpha_1 + \mathbb{Z}6\alpha_2 + \mathbb{Z}2\alpha_3 + \mathbb{Z}4\alpha_4.
\end{aligned}$$

Now,

$$\mathbb{Z}6\alpha_1 + \mathbb{Z}6\alpha_2 + \mathbb{Z}12\alpha_3 + \mathbb{Z}12\alpha_4 \subset [M]_W.$$

Moreover, $\mathbb{Z}6\alpha_1 + \mathbb{Z}6\alpha_2 + \mathbb{Z}12\alpha_3 + \mathbb{Z}12\alpha_4$ is W -stable (as can be easily seen). Conversely, take $\mu \in [M]_W$. Then, $w\mu \in M$ for any $w \in W$. Since

$$s_i w\mu = w\mu - (w\mu)(\alpha_i^\vee)\alpha_i;$$

for any $\mu \in [M]_W$ and any $w \in W$,

$$w\mu(\alpha_2^\vee) \in 6\mathbb{Z} \text{ and } w\mu(\alpha_4^\vee) \in 4\mathbb{Z}.$$

Since $\alpha_1^\vee \in W.\alpha_2^\vee$ (cf [H, §10.4, Lemma c]), we get

$$\mu(\alpha_1^\vee) \in 6\mathbb{Z}, \mu(\alpha_2^\vee) \in 6\mathbb{Z} \text{ and } \mu(\alpha_4^\vee) \in 4\mathbb{Z}.$$

Take $\mu = n_1 2\alpha_1 + n_2 6\alpha_2 + n_3 2\alpha_3 + n_4 4\alpha_4 \in [M]_W$. Then, from the above, we get

$$4n_1 - 6n_2 \in 6\mathbb{Z}, -2n_1 + 12n_2 - 2n_3 \in 6\mathbb{Z} \text{ and } -2n_3 + 8n_4 \in 4\mathbb{Z}.$$

This gives that $n_1 \in 3\mathbb{Z}$ and $n_3 \in 6\mathbb{Z}$. Thus, $\mu \in \mathbb{Z}6\alpha_1 + \mathbb{Z}6\alpha_2 + \mathbb{Z}12\alpha_3 + \mathbb{Z}4\alpha_4$. Considering $s_3(\mathbb{Z}6\alpha_1 + \mathbb{Z}6\alpha_2 + \mathbb{Z}12\alpha_3 + \mathbb{Z}4\alpha_4)$, we see that

$$[\mathbb{Z}6\alpha_1 + \mathbb{Z}6\alpha_2 + \mathbb{Z}12\alpha_3 + \mathbb{Z}4\alpha_4]_W \subset \mathbb{Z}6\alpha_1 + \mathbb{Z}6\alpha_2 + \mathbb{Z}12\alpha_3 + \mathbb{Z}12\alpha_4.$$

Thus, we conclude that

$$[M]_W = \mathbb{Z}6\alpha_1 + \mathbb{Z}6\alpha_2 + \mathbb{Z}12\alpha_3 + \mathbb{Z}12\alpha_4.$$

This completes the proof of the theorem in the case of F_4 .

(g) E_6 : By the chart and the case of A_ℓ ,

$$\begin{aligned} L(E_6; \alpha_1, \dots, \alpha_6) &= [(\mathbb{Z}\theta + L(A_5; \alpha_1, \alpha_3, \alpha_4, \alpha_5, \alpha_6)) \cap \\ &\quad (\mathbb{Z}\alpha_1 + L(A_5; -\theta, \alpha_2, \alpha_4, \alpha_5, \alpha_6)) \cap \\ &\quad (L(A_2; \alpha_1, \alpha_3) + L(A_2; -\theta, \alpha_2) + L(A_2; \alpha_5, \alpha_6)) \cap \\ &\quad (L(A_5; \alpha_1, \alpha_3, \alpha_4, \alpha_2, -\theta) + \mathbb{Z}\alpha_6)]_W \\ &= [(\mathbb{Z}\theta + \mathbb{Z}\alpha_1 + \mathbb{Z}\alpha_3 + \mathbb{Z}\alpha_4 + \mathbb{Z}\alpha_5 + \mathbb{Z}\alpha_6) \cap \\ &\quad (\mathbb{Z}\alpha_1 + \mathbb{Z}\theta + \mathbb{Z}\alpha_2 + \mathbb{Z}\alpha_4 + \mathbb{Z}\alpha_5 + \mathbb{Z}\alpha_6) \cap \\ &\quad (\mathbb{Z}\alpha_1 + \mathbb{Z}\alpha_3 + \mathbb{Z}\theta + \mathbb{Z}\alpha_2 + \mathbb{Z}\alpha_5 + \mathbb{Z}\alpha_6) \cap \\ &\quad (\mathbb{Z}\alpha_1 + \mathbb{Z}\alpha_3 + \mathbb{Z}\alpha_4 + \mathbb{Z}\alpha_2 + \mathbb{Z}\theta + \mathbb{Z}\alpha_6)]_W \\ &= [(\mathbb{Z}\alpha_1 + \mathbb{Z}2\alpha_2 + \mathbb{Z}\alpha_3 + \mathbb{Z}\alpha_4 + \mathbb{Z}\alpha_5 + \mathbb{Z}\alpha_6) \cap \\ &\quad (\mathbb{Z}\alpha_1 + \mathbb{Z}\alpha_2 + \mathbb{Z}2\alpha_3 + \mathbb{Z}\alpha_4 + \mathbb{Z}\alpha_5 + \mathbb{Z}\alpha_6) \cap \\ &\quad (\mathbb{Z}\alpha_1 + \mathbb{Z}\alpha_2 + \mathbb{Z}\alpha_3 + \mathbb{Z}3\alpha_4 + \mathbb{Z}\alpha_5 + \mathbb{Z}\alpha_6) \cap \\ &\quad (\mathbb{Z}\alpha_1 + \mathbb{Z}\alpha_2 + \mathbb{Z}\alpha_3 + \mathbb{Z}\alpha_4 + \mathbb{Z}2\alpha_5 + \mathbb{Z}\alpha_6)]_W \\ &= [M_6]_W, \end{aligned}$$

where $M_6 := \mathbb{Z}\alpha_1 + \mathbb{Z}2\alpha_2 + \mathbb{Z}2\alpha_3 + \mathbb{Z}3\alpha_4 + \mathbb{Z}2\alpha_5 + \mathbb{Z}\alpha_6$.

Clearly, $6\Lambda \subset M_6$ and since Λ is W -stable,

$$(1) \quad 6\Lambda \subset [M_6]_W.$$

Conversely, take $\mu \in [M_6]_W$. Then, for any $w \in W$, $w\mu \in M_6$. Since $s_i(w\mu) = w\mu - (w\mu)(\alpha_i^\vee)\alpha_i$; for any $\mu \in [M]_W$, and any $w \in W$, $w\mu(\alpha_2^\vee) \in 2\mathbb{Z}$ and $w\mu(\alpha_4^\vee) \in 3\mathbb{Z}$. Since E_6 is simplylaced, the Weyl group W acts transitively on the coroots [H, §10.4, Lemma C]. Thus, $\mu(\alpha_i^\vee) \in 2\mathbb{Z} \cap 3\mathbb{Z} = 6\mathbb{Z}$ for all the simple coroots $\alpha_i^\vee$. This proves that $\mu \in 6\Lambda$, i.e.,

$$(2) \quad [M_6]_W \subset 6\Lambda.$$

Comparing (1) and (2) we get $[M_6]_W = 6\Lambda$. This proves the theorem in the case of E_6 .

h) E_7 : By the chart and the result for the cases of A_ℓ, D_ℓ , we get

$$\begin{aligned} L(E_7; \alpha_1, \dots, \alpha_7) &= [(\mathbb{Z}\theta + L(D_6; \alpha_7, \alpha_6, \dots, \alpha_2)) \cap \\ &\quad L(A_7; -\theta, \alpha_1, \alpha_3, \alpha_4, \dots, \alpha_7) \cap \\ &\quad (L(A_2; -\theta, \alpha_1) + L(A_5; \alpha_2, \alpha_4, \alpha_5, \alpha_6, \alpha_7)) \cap \\ &\quad (L(A_5; -\theta, \alpha_1, \alpha_3, \alpha_4, \alpha_2) + L(A_2; \alpha_6, \alpha_7)) \cap \\ &\quad (L(D_6; -\theta, \alpha_1, \alpha_3, \alpha_4, \alpha_5, \alpha_2) + \mathbb{Z}\alpha_\ell)]_W \\ &= [(\mathbb{Z}\theta + L'(\alpha_7, \alpha_6, \dots, \alpha_2)) \cap \\ &\quad (\mathbb{Z}\theta + \mathbb{Z}\alpha_1 + \mathbb{Z}\alpha_3 + \mathbb{Z}\alpha_4 + \dots + \mathbb{Z}\alpha_7) \cap \\ &\quad (\mathbb{Z}\theta + \mathbb{Z}\alpha_1 + \mathbb{Z}\alpha_2 + \mathbb{Z}\alpha_4 + \dots + \mathbb{Z}\alpha_7) \cap \\ &\quad (\mathbb{Z}\theta + \mathbb{Z}\alpha_1 + \dots + \mathbb{Z}\alpha_4 + \mathbb{Z}\alpha_6 + \mathbb{Z}\alpha_7) \cap \\ &\quad (L'(-\theta, \alpha_1, \alpha_3, \alpha_4, \alpha_5, \alpha_2) + \mathbb{Z}\alpha_7)]_W \\ &= [M_7]_W, \end{aligned}$$

where $M_7 := \mathbb{Z}4\alpha_1 + \mathbb{Z}2\alpha_2 + \mathbb{Z}6\alpha_3 + \mathbb{Z}2\alpha_4 + \mathbb{Z}6\alpha_5 + \mathbb{Z}4\alpha_6 + \mathbb{Z}2\alpha_7$.

Clearly, $12\Lambda \subset M_7$ and since Λ is W -stable, $12\Lambda \subset [M_7]_W$.

For any $\mu \in [M_7]_W$, by considering $(w\mu)(\alpha_1^\vee)$ and $(w\mu)(\alpha_3^\vee)$ as in the proof of the theorem for the case of E_6 , we get that $\mu(\alpha_i^\vee) \in 4\mathbb{Z} \cap 6\mathbb{Z} = 12\mathbb{Z}$ for all the simple coroots $\alpha_i^\vee$. Thus,

$$[M_7]_W \subset 12\Lambda \text{ and hence } [M_7]_W = 12\Lambda.$$

This takes care of the case of E_7 .

Finally, we come to the following:

(i) E_8 : By the chart and the theorem for E_6, E_7, D_8 and A_ℓ , we get (denoting the weight lattice of E_i by $\Lambda(E_i)$).

$$\begin{aligned}
L(E_8; \alpha_1, \dots, \alpha_8) &= [L(D_8; -\theta, \alpha_8, \alpha_7, \dots, \alpha_2) \cap \\
&\quad L(A_8; \alpha_1, \alpha_3, \alpha_4, \dots, \alpha_8, -\theta) \cap \\
&\quad (L(A_4; \alpha_1, \alpha_3, \alpha_4, \alpha_2) + L(A_4; \alpha_6, \alpha_7, \alpha_8, -\theta)) \cap \\
&\quad (L(E_6; \alpha_1, \dots, \alpha_6) + L(A_2; \alpha_8, -\theta)) \cap \\
&\quad (L(E_7; \alpha_1, \dots, \alpha_7) + \mathbb{Z}\theta)]_W \\
&= [L'(\theta, \alpha_8, \alpha_7, \dots, \alpha_2) \cap \\
&\quad (\mathbb{Z}\alpha_1 + \mathbb{Z}\alpha_3 + \mathbb{Z}\alpha_4 + \dots + \mathbb{Z}\alpha_8 + \mathbb{Z}\theta) \cap \\
&\quad (\mathbb{Z}\alpha_1 + \dots + \mathbb{Z}\alpha_4 + \mathbb{Z}\alpha_6 + \mathbb{Z}\alpha_7 + \mathbb{Z}\alpha_8 + \mathbb{Z}\theta) \cap \\
&\quad (6\Lambda(E_6) + \mathbb{Z}\alpha_8 + \mathbb{Z}\theta) \cap (12\Lambda(E_7) + \mathbb{Z}\theta)]_W \\
&= [M'_8 \cap (6\Lambda(E_6) + \mathbb{Z}\alpha_8 + \mathbb{Z}\theta) \cap (12\Lambda(E_7) + \mathbb{Z}\theta)]_W,
\end{aligned}$$

where

$$M'_8 := \{\mathbb{Z}4\alpha_1 + n3\alpha_2 + m\alpha_3 + \mathbb{Z}2\alpha_4 + \mathbb{Z}10\alpha_5 + \mathbb{Z}2\alpha_6 + \mathbb{Z}2\alpha_7 + \mathbb{Z}2\alpha_8 : n, m \in \mathbb{Z} \text{ and } n+m \text{ is even}\}.$$

But the coefficient of α_3 in any element of $6\Lambda(E_6) + \mathbb{Z}\alpha_8 + \mathbb{Z}\theta$ is even. Thus,

$$\begin{aligned}
[M'_8 \cap (6\Lambda(E_6) + \mathbb{Z}\alpha_8 + \mathbb{Z}\theta) \cap (12\Lambda(E_7) + \mathbb{Z}\theta)]_W &= \\
[M_8 \cap (6\Lambda(E_6) + \mathbb{Z}\alpha_8 + \mathbb{Z}\theta) \cap (12\Lambda(E_7) + \mathbb{Z}\theta)]_W,
\end{aligned}$$

where

$$M_8 := \mathbb{Z}4\alpha_1 + \mathbb{Z}6\alpha_2 + \mathbb{Z}2\alpha_3 + \mathbb{Z}2\alpha_4 + \mathbb{Z}10\alpha_5 + \mathbb{Z}2\alpha_6 + \mathbb{Z}2\alpha_7 + \mathbb{Z}2\alpha_8.$$

For any $\mu \in [M_8]_W$, by considering $(w\mu)(\alpha_1^\vee), (w\mu)(\alpha_2^\vee)$ and $(w\mu)(\alpha_5^\vee)$ as in the proof of the theorem for the case of E_6 , we get

$$(3) \quad [M_8]_W \subset 60\Lambda(E_8).$$

Conversely, since $\Lambda(E_8) = Q(E_8)$, we get that

$$(4) \quad 60\Lambda(E_8) \subset M_8 \cap (6\Lambda(E_6) + \mathbb{Z}\alpha_8 + \mathbb{Z}\theta) \cap (12\Lambda(E_7) + \mathbb{Z}\theta),$$

and hence

$$60\Lambda(E_8) \subset [M_8 \cap (6\Lambda(E_6) + \mathbb{Z}\alpha_8 + \mathbb{Z}\theta) \cap (12\Lambda(E_7) + \mathbb{Z}\theta)]_W.$$

To prove (4), it suffices to show that

$$(5) \quad 60\alpha_7 \in 6\Lambda(E_6) + \mathbb{Z}\alpha_8 + \mathbb{Z}\theta$$

and

$$(6) \quad 60\alpha_8 \in 12\Lambda(E_7) + \mathbb{Z}\theta.$$

To prove (5), observe that

$$\begin{aligned} 60\alpha_7 &= 20\theta - 40\alpha_8 - 20(2\alpha_1 + 3\alpha_2 + 4\alpha_3 + 6\alpha_4 + 5\alpha_5 + 4\alpha_6) \\ &= 20\theta - 40\alpha_8 - 60\omega_6(E_6), \end{aligned}$$

where $\omega_6(E_6)$ is the sixth fundamental weight of E_6 . Similarly, to prove (6), observe that

$$\begin{aligned} 60\alpha_8 &= 30\theta - 30(2\alpha_1 + 3\alpha_2 + 4\alpha_3 + 6\alpha_4 + 5\alpha_5 + 4\alpha_6 + 3\alpha_7) \\ &= 30\theta - 60\omega_7(E_7). \end{aligned}$$

This proves (5) and (6) and thus (4). Combining (3) and (4), we get that

$$[M_8 \cap (6\Lambda(E_6) + \mathbb{Z}\alpha_8 + \mathbb{Z}\theta) \cap (12\Lambda(E_7) + \mathbb{Z}\theta)]_W = 60\Lambda(E_8) = 60Q.$$

This proves the theorem for E_8 and hence the theorem is completely proved. $\square$

Remark 3.11 *Theorem 3.10 (a) is obtained earlier by B. Howard [Ho].*

References

- [B] N. Bourbaki: *Groupes et Algèbres de Lie, Chap. 4-6*, Masson, Paris (1981).
- [DN] J.-M. Drezet and M.S. Narasimhan: Groupe de Picard des variétés de modules de fibrés semi-stables sur les courbes algébriques, *Invent. Math.* **97** (1989), 53–94.
- [H] J.E. Humphreys: *Introduction to Lie Algebras and Representation Theory*, GTM vol. **9**, Springer-Verlag (1972).
- [Ho] B.J. Howard: Matroids and geometric invariant theory of torus actions on flag spaces, Preprint (2005).
- [MFK] D. Mumford, J. Fogarty and F. Kirwan: *Geometric Invariant Theory*, Third Edition, Ergebnisse der Mathematik und ihrer Grenzgebiete vol. **34**, Springer (1994).
- [W] J. Wolf: *Spaces of constant curvature*, Third Edition, Publish or Perish, Inc. (1974).

Address: Department of Mathematics, UNC at Chapel Hill, Chapel Hill, NC 27599-3250, USA (email: shrawan@email.unc.edu)